

AN ASCA STUDY OF THE SUPERNOVA REMNANT G39.2–0.3

Ilana M. Harrus^{1,2} and Patrick O. Slane^{1,3}

Received August 1998; accepted November 1998

For submission to *The Astrophysical Journal*

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138-1516

² imh@head-cfa.harvard.edu

³ slane@cfa.harvard.edu

ABSTRACT

We present the analysis of archival data from the *Advanced Satellite for Cosmology and Astrophysics* (*ASCA*) of the supernova remnant (SNR) G39.2–0.3. G39.2–0.3 has been sometimes characterized as a shell-like remnant in the literature but our high-energy imaging and spectral analysis show the unambiguous composite nature of the remnant. We find that part of the contribution to G39.2–0.3’s X-ray emission is distinctly non-thermal, best described by a power law with a photon index ($2.53^{+0.34}_{-0.27}$). The region of emission is consistent with a point source (extension consistent with the point spread function of the detector at the off-axis angle of the observation) and is roughly defined by a circle of radius $\sim 4'$. A second contribution comes from a thermal component which contributes flux primarily at low energies. Despite the absence of any pulsed emission detected from the compact source, we argue that the emission is most probably due to a rotating compact object that is powering the detected synchrotron nebula within the SNR.

Subject headings: radiation mechanisms: non-thermal — supernova remnants: individual: G39.2–0.3 — X-rays: ISM

1. Introduction

The standard theory of core collapse of a massive star predicts the formation of a compact object (neutron star or black hole) as the normal by-product of such a supernova explosion. Indeed the most famous example of such an object is the Crab nebula whose central pulsar has been extensively studied for nearly three decades, studies which have led to the development of our understanding of the physics of the interaction of the compact object and its surroundings. In addition to the 33 msec-pulsar studies, the spectral analysis of the Crab Nebula and its featureless power-law spectrum has brought to light the emission mechanisms at work in the synchrotron nebula. Considering that nearly all Type Ib/c or Type II supernovae result in the formation of a neutron star, it has been suggested that the small number of confirmed pulsar/SNR associations (less than a dozen) is a puzzle (Narayan & Schaudt 1988). Some authors argue that this can be well explained by detection limits of faint pulsars, statistical fluctuations and life time differences between SNRs and pulsars (Gaensler & Johnston 1995). It remains true, however, that the class of pulsars for which the emission beam does not intersect with the Earth is lost to pulsed emission searches. Recent developments in X-ray observations have made it possible to circumvent this problem. Broadband X-ray observations have provided a tool to confirm the presence of a compact object by detecting the associated synchrotron nebula or plerion. This method does not suffer from the beaming effect bias and gives direct information on the current energy output of the central powering object. With the advance of the high sensitivity broadband X-ray imaging and spectral instruments onboard *ASCA*, the study of this class of objects has made great progress.

The classification of G39.2–0.3 has always been somewhat unclear. The remnant was classified as a Crab-like SNR despite the fact that its spectral index was more typical of a shell-type object (Caswell et al 1975; Caswell et al 1982) both because the maximum

surface brightness at 1.4GHz is located at the center of the remnant, and in view of the large polarization measured across the remnant. Polarization is measured both at 1400 and 500 MHz (Caswell et al 1982). It is distributed over the remnant with a peak of 10% polarization in the western and central parts and 50% in the eastern part.

Becker & Helfand (1987) argued that the enhanced emission was most probably due to a projection effect as opposed to the emission of a plerionic component. In view of the shell-like spectral index and the lack of any detected spectral variation across the remnant, the authors concluded that G39.2–0.3 was most probably a shell-like remnant.

Another observation of G39.2–0.3 made by Patnaik et al (1990) using the Very Large Array (VLA), confirmed the typical shell-like integrated spectral index of the remnant (0.42 ± 0.02). G39.2–0.3 proved to be more surprising: Anderson & Rudnick (1993) found that although the average index of the remnant was indeed around the value found by Patnaik et al (1990), G39.2–0.3 showed evidence of spatial variation of the spectral index, from ($0.58^{+0.04}_{-0.05}$) at the edge to a Crab-like index of ($0.39^{+0.10}_{-0.05}$) at the center. In the high-energy band, G39.2–0.3 was observed by the *Einstein* Imaging Proportional Counter (IPC), but the total observation time was small (2568 sec). Although clearly detected, the total number of photons (80 ± 10 photons) was insufficient to differentiate between a thermal and a non-thermal spectrum (Becker & Helfand 1987). No on-axis pointing of the object was made with either instrument onboard *ROSAT* (the Position Sensitive Proportional Counter or the High Resolution Imager).

In view of neutral-hydrogen absorption measurements (Caswell et al 1975) a minimum distance between 7.7 and 11.3 kpc is generally assumed. Becker & Helfand (1987) estimated that the hydrogen column-density toward the remnant is 2×10^{22} atoms cm^{-2} based on 21cm absorption measurements with the VLA and concurred on the estimate of the distance.

Presented here are the results and interpretations of archival *ASCA* X-ray data on the SNR G39.2–0.3. We present in §2 our data extraction methods and spatial, spectral, and

timing analyses. In §3 we discuss the implications of our results. The final section of the article is a summary of our principal conclusions.

2. Data Reduction and Analysis

We retrieved the *ASCA* data for G39.2–0.3 directly from the HEASARC (High Energy Astrophysics Science Archive Research Center) public database. Data provided with standard screening process were used for the analysis (see http://adfwww.gsfc.nasa.gov/asca/processing_doc/proc/processing.html for more information on that screening process). No special configuration was applied to the time-recording data bits and the time resolution is the nominal one: 0.5 sec in medium-bit-rate mode and 0.0625 sec in high-bit-rate mode for the GIS detector (the SIS is not used for timing analysis).

2.1. Spatial Analysis

We have generated exposure-corrected, background-subtracted images of the GIS (GIS 2 and GIS 3 combined) and SIS (SIS 0 and SIS 1 combined) data in soft (below 3.0 keV) and hard (above 3.0 keV) energy bands for the 40 ksec observation. The background for the spatial analysis was determined from the weighted average of several nominally blank fields from high Galactic latitude observations with data selection criteria matched to those used for the SNR data. Exposure maps were generated from the off-axis effective-area calibrations, weighted by the appropriate observation time. Events from regions of the merged exposure map with less than 10% of the maximum exposure were ignored. Merged images of the source data, background, and exposure were smoothed with a Gaussian of standard deviation, $\sigma = 45''$. We subtracted smoothed background maps from the data maps and divided by the corresponding exposure map.

Fig. 1 shows the results of this procedure for both the SIS and the GIS detectors. The high-energy emission, both in the GIS and the SIS, appears to be more centrally enhanced than the low-energy one. At high energy the maximum emission, identical for both SIS and GIS, is located at $19^{\text{h}}04^{\text{m}}4.8^{\text{s}}$, $05^{\circ}27'11.2''$ (J2000). The center of emission is slightly different in the soft and hard band (we measure a $22''$ offset mainly along the North-South axis). Considering the off-axis angle of the X-ray pointing (the observation was made with an average off-axis angle of $7'15''$) and the broad point-spread function of the detector, such an apparent offset may be purely associated with systematic overcorrection of vignetting effects. Similarly, we are unable to definitely establish the point-like or extended nature of the high-energy emission. The high-energy X-ray emission region is compatible with a point source (we used a circle of radius $\simeq 4'$ to extract events for both the SIS and the GIS).

Fig. 2 shows both a 20cm and a 6cm polarized-intensity radio map (data from Patnaik et al 1990) superimposed with the SIS contours in the high-energy band (from 3.0 to 10.0 keV). We note the strong positional correlation between the maximum of the polarized-emission and the center of the high-energy X-ray profile.

2.2. Spectral Analysis

We extracted X-ray spectra from circular regions centered at $19^{\text{h}}04^{\text{m}}2.1^{\text{s}}$, $05^{\circ}26'54''$ (J2000) for both GIS and SIS, using a radius of $4'17''$ in the GIS and only $4'$ in the SIS because of the CCD-chip boundaries.

After the standard data selection criteria were applied, the spatial cuts described above lead to combined counts rates of 0.057 ± 0.003 cnt sec $^{-1}$ and 0.068 ± 0.002 cnt sec $^{-1}$ for the SIS and the GIS respectively.

We first fitted the SIS and GIS separately and then jointly after making sure that the combined fits were compatible with the separated fit. All results described below were

obtained with a simultaneous fit of all four sets of data. As seen in the spatial analysis, the emission from the remnant is localized and centrally peaked, so it was possible to extract background spectra from the same field of view both for the SIS and the GIS. This method has the advantage of eliminating a possible high-energy contamination from the Galactic plane.

The models applied to fit the emission profile of the remnant reflect the physics of the phenomena which could be contributing to the X-ray output of the remnant. If the X-ray emission is coming from the interaction of a blast wave with the interstellar medium (ISM) the emission is expected to be thermal. Several models are used in X-ray astrophysics to describe such plasmas. In the first part of our analysis, we have modeled the thermal emission with a collisional equilibrium ionization (CEI) model (the so-called “mekal” model - Mewe, Gronenschild & van den Oord 1985; Mewe, Lemen & van den Oord 1986; Kaastra 1992) which is available in the version 10.00 of XSPEC, the X-ray spectral analysis package used throughout this analysis. (The other collisional equilibrium model available, Raymond & Smith 1977, leads to similar results). Unless explicitly stated, we have kept the elemental abundances at their nominal values as defined by Anders & Grevesse (1989).

Absorption along the line of sight was taken into account with an equivalent column density of hydrogen, N_{H} , using the cross-sections and abundances from Morrison & McCammon (1983). In all of the following analysis, the gain offsets of both GIS detectors are linked together and allowed to vary freely. A gain shift around -3% is measured in both detectors, consistent with the results from the calibration data analysis done by the *ASCA*-GIS team (see http://heasarc.gsfc.nasa.gov/docs/asca/cal_probs.html for more information on calibration issues). We link the normalizations of SIS 0 and SIS 1 spectra (and similarly for GIS 2 and GIS 3) while allowing for small variations between the SIS and GIS. We checked that the results for each detector (SIS and GIS) are consistent with each other at the 1σ level, considering that due to the CCD boundaries, some flux is consistently

missed in the SIS fit normalization. As a result, all fluxes quoted hereafter are derived from the GIS results.

We fitted a pure thermal (CEI model) description of the emission and obtained marginally good fit with a χ^2 of 422.3 (and a reduced $\chi_r^2 = 1.77$). The other single model we tried on the four data sets, is a pure non-thermal model. In this case, we found a worse χ^2 of 492.0 with an associated $\chi_r^2=2.07$. In both cases the associated column density is above 3×10^{22} atoms cm^{-2} consistent with the derived values from the IPC analysis (Becker & Helfand 1987). A combined fit including both thermal and non-thermal contributions leads to a drop of more than 200 in χ^2 (to a value of 241.6) and a $\chi_r^2=1.02$. The hydrogen column density is high, $(4.70 \pm 0.30) \times 10^{22}$ atoms cm^{-2} , as expected from the distance estimate and the location of the remnant, and the temperature is $kT = 0.61 \pm 0.10$ keV. The power-law fit leads to a photon index of $2.08^{+0.46}_{-0.25}$. We also tried to fit the high-energy part of the spectrum with a bremsstrahlung model. In this case, the temperature associated is high ($4.0^{+2.0}_{-1.0}$ keV). The column density is consistent to the one found with the previous model ($4.71^{+0.35}_{-0.33} \times 10^{22}$ atoms cm^{-2}) and so is the temperature of the low-energy component ($kT = 0.51^{+0.08}_{-0.06}$ keV). The χ^2 is worse ($\chi_r^2=1.16$) than the one found with the previous power-law/thermal mixed model and we favor the power-law interpretation of the high-energy component.

We note that the column density is high relative to that determined from HI absorption measurements. Such discrepancies are not uncommon: Arabadjis & Bregman (1998) show that, at low galactic latitudes, X-ray derived values are roughly double that of the ones deduced from 21 cm measurements, presumably indicating the presence of additional molecular material.

We have also examined the effects of non-equilibrium ionization (NEI) on our previous results. Non equilibrium effects arise when the ions are not instantaneously ionized to their equilibrium temperature behind the shock front. Rather, the timescale for attaining full equilibrium ionization is comparable to the remnant dynamical timescale. To incorporate these effects into the model of SNR spectral emissivity, we use here the matrix inversion

method developed by Hughes & Helfand (1985). The ionization state depends on the product of electron density and age and we define the ionization timescale as $\tau_i \equiv n_e t$. As in the CEI model, we have kept all elements to their nominal abundances (Anders & Grevesse 1989). From the results obtained in the previous analysis, we can estimate an upper limit for the expected value of the ionization timescale. We find an upper limit for $n_e t$ between 6780 and 15606 yrs cm⁻³. As in the CEI analysis, a single model fit cannot account for the total spectra. In the case of a single NEI model, we find $\chi^2 = 287.80$ for 232 degrees of freedom larger than the best χ^2 of the complete analysis but much better than the one obtained in the equilibrium case alone. Upon adding a power-law component we find a $\chi^2 = 238.11$ (for 235 degrees of freedom). The associated column density is $(4.65^{+0.11}_{-0.26}) \times 10^{22}$ atoms cm⁻². The temperature ($kT = 0.62^{+0.10}_{-0.03}$ keV) is in complete agreement with the one found in the CEI model and the ionization timescale (3810^{+2095}_{-1510} yrs cm⁻³) is consistent with the derived upper limit given above. The power-law model gives a slightly steeper photon index of $2.53^{+0.34}_{-0.27}$. This good agreement between the equilibrium and non-equilibrium models argues against a very young remnant for which non-equilibrium effects would be very strong, a fact that is confirmed by the age estimate derived in the following section. We have searched for variation in elemental abundances (in particular for Magnesium, Silicon and Sulfur) and found none. Fig. 3 shows the data and the best-fit power-law plus NEI thermal model. The thermal component is plotted separately in the insert to the figure. It is clear from the figure alone that the thermal contribution is prevalent at low energy while it is contributing only a small amount above 3.0 keV. We argue that the spectral data support the interpretation that the X-ray emission from G39.2–0.3 consists of non-thermal synchrotron emission from a central plerion in addition to a thermal component arising from the interaction of the blast wave with the ISM. Results from the spectral fits are tabulated in Table 1.

2.3. Timing Analysis

As noted in our study of MSH 11–62 (Harrus et al 1998) it is often the case that no pulse is detected from SNRs which present unambiguous evidence for a compact object; beaming effects, for example, may result in a cone of emission which does not intersect the Earth. This beaming effect presumably accounts for the relatively low number of SNRs for which an associated radio pulsar has been unambiguously identified. In our timing analysis of G39.2–0.3, we have extracted data from the central part of the emission and restricted ourselves to energies above 3 keV in order to suppress the contribution from the thermal spectral component associated with the remnant. We searched for periodicity in intervals of 0.0625 sec (the best time resolution of our data) up to 20 sec. No significant deviation from a random distribution is found. To insure the validity of our method, we have extracted from the HEASARC archives the data for an SNR in which a similar analysis lead to the discovery of a pulse emission. Identical cuts (both in position and in energy) are applied to Kes 73 data and the same search leads to the detection (at more than 10σ level) of the pulsar located at the center of this remnant (Vasisht & Gotthelf 1997). On the total 2349 events extracted for the G39.2–0.3 timing analysis, we estimate a background from the thermal component around 900 events and we derived a pulsed fraction upper limit between 10% and 25% (depending on the pulse shape) at the 90% confidence level.

3. Discussion

Using a distance compatible with the minimum distance to G39.2–0.3 between 7.7 and 11.3 kpc (Caswell et al 1975), the supernova remnant defines an X-ray emitting volume of $V \simeq 1.94 \times 10^{59} f D_{10}^3 \theta_4^3 \text{ cm}^{-3}$, where f is the volume filling factor of the emitting gas within the SNR, D_{10} is the distance to the remnant in units of 10 kpc, and θ_4 the angular radius in units of $4'$. From this estimate, the combined result from the thermal component (see Table 1), and a ratio n_e/n_H of 1.08 (extracted from our NEI analysis), we deduce a

hydrogen number density of $n_{\text{H}} = (0.80^{+0.20}_{-0.17}) D_{10}^{-1/2} \theta_4^{-3/2} f^{-1/2} \text{ cm}^{-3}$. The mass of X-ray emitting plasma is $M_X = 175 D_{10}^{5/2} f^{1/2} \theta_4^{3/2} M_{\odot}$, which for the distance range mentioned above and combined with the errors from the spectral analysis, translates to a mass M_X between 40 and 300 $M_{\odot}$. Considering that the distances used are already a lower bound of the possible distance to the remnant, we argue that this result is consistent with the remnant being in its Sedov-Taylor phase of evolution (Taylor 1950; Sedov 1959). In this case, an estimate on the age of the remnant can be made using the temperature obtained in our spectral analysis. This temperature, $\langle T \rangle$, is the emission-measure-weighted average electron temperature, which is proportional to the shock temperature $\langle T \rangle = 1.27 T_S$ (assuming that electrons and ions have the same temperature). The Sedov age relationship is $t \approx 480 \text{ yrs } (R/1 \text{ pc}) (\langle T \rangle / 1 \text{ keV})^{-1/2} \approx (7100^{+1150}_{-2020}) \text{ yr}$ (the errors bars reflect both the 1σ errors on the measured temperature and the uncertainties on the distance). The preshock ISM number density is $n_0 = \rho_0 m_{\text{H}} = (0.24^{+0.07}_{-0.05}) D_{10}^{-1/2} \theta_4^{-3/2} f^{-1/2} \text{ cm}^{-3}$, which leads to $n_0 = (0.49^{+0.14}_{-0.10}) D_{10}^{-1/2} \theta_4^{-3/2} \text{ cm}^{-3}$ for a filling factor of 0.25 (as expected in the case of a pure Taylor-Sedov solution). With the preceding numerical values we estimate the supernova explosion energy from the Taylor-Sedov relations to be $E = (3.5^{+1.5}_{-0.8}) \times 10^{50} D_{10}^{5/2} \theta_4^{3/2} f^{-1/2} \text{ ergs}$ which is within the range of typical values.

Assuming temperature equilibrium between electrons and ions, we can compute the pressure at the shock front $P_s \sim 1.93 n_e k T_S \sim (1.29^{+0.61}_{-0.33}) \times 10^{-9} D_{10}^{-1/2} f^{-1/2} \text{ dyne cm}^{-2}$. For a Taylor-Sedov temperature and density distribution, the central pressure is $P_0 \sim 0.31 P_s = 8 \times 10^{-10} \text{ dyne cm}^{-2}$ for a filling factor of 0.25. This value of the internal thermal pressure leads to an equipartition magnetic field value $B = \sqrt{8\pi P_0} = 100^{+20}_{-10} D_{10}^{-1/4} f^{-1/4} \mu\text{G}$. This estimate of the magnetic field allows us to estimate the break frequency ν_{B} , the frequency at which synchrotron losses begin to dominate. We have

$$B = 14.6 (\nu_{\text{B}} / 10^6 \text{ GHz})^{-1/3} (t / 10^4 \text{ yrs})^{-2/3} \mu\text{G}. \quad (1)$$

(Ginzburg & Syrovatskii 1965 eqn. 5.36). Here t is the age of the nebula for which we use the value derived in our study of the thermal emission. For the results mentioned above and including the uncertainties on the distance and the magnetic field, we find a break frequency between 3350 and $1.25 \times 10^4 f^{3/4}$ GHz (to be compared to the Crab break frequency $\nu_{\text{break}} = 10^4$ GHz)(see Table 2).

The combined existing radio and X-ray data can provide another estimate of the range of possible break frequencies. Using the spectral index ($0.39^{+0.10}_{-0.05}$) and the measured radio flux from the central region of G39.2–0.3 (0.20 mJy arcsec $^{-2}$) (Anderson & Rudnick 1993), we found a range of frequencies ranging from 1.4×10^5 to 3×10^6 GHz. To reconcile the apparent discrepancy between the two ranges, one must note that the actual plerion contribution to the radio flux is smaller than the total observed flux (which includes contribution from the shell) and that the spectral index is probably overestimated as well. Both values affect the value of the break frequency range quoted above. As an example, we found that for a spectral index of 0.20 and a filling factor of 0.5, the two ranges of frequency are overlapping.

We have argued above that the nonthermal emission is produced by a synchrotron nebula surrounding a spinning neutron star, or pulsar, created in the supernova explosion. Using the measured X-ray luminosity associated with the non-thermal component, $L_X^\alpha = 2.28 \times 10^{35} D_{10}^2$ ergs s $^{-1}$ (0.2 – 4.0 keV) along with the empirical relationship between L_X^α and the pulsar’s spin-down energy $\dot{E}$ (Seward & Wang 1988), we infer a $\dot{E} \sim 2.4 \times 10^{37} D_{10}^{1.44}$ ergs s $^{-1}$ for the putative pulsar in G39.2–0.3. Using this value, an assumed breaking index of 3, and the age deduced from the Sedov analysis, we derive an initial spin period range of 24–36 ms for surface magnetic field values between 10^{12} and 7×10^{12} G. From this initial spin period, one can deduce the current-epoch period in the range of 25 to 70 ms.

SNRs with similar characteristics have been found with *ASCA*. The class includes

MSH 11–62 (Harrus, Hughes & Slane 1998) and Kes 75 (Blanton & Helfand 1996) which have similar values for the age (depending upon distance), photon index, and temperature. In fact among the objects which belong to the class of SNRs for which *ASCA* has revealed a high-energy emitting center, G39.2–0.3 appears to be rather average.

4. Summary

We have presented the results of *ASCA* X-ray spectral and spatial studies of the SNR G39.2–0.3, a middle-aged remnant (about 7000 year old) in its Taylor-Sedov phase of evolution. In our model, its X-ray emission consists of 1) a thermal emission from the blast wave interacting with the ISM and 2) a non-thermal synchrotron emission due to high-energy electrons surrounding the compact object. We can develop an acceptable description for the synchrotron nebula with an associated spin-down power of the undetected and central neutron star of $\dot{E} = 2.4 \times 10^{37} \text{ erg s}^{-1}$.

This research has made use of data obtained from the High Energy Astrophysics Science Archive Research Center (HEASARC), provided by NASA’s Goddard Space Flight Center. This research was supported in part by NASA under grants NAG5-3486 and contract NAS8-39073. The authors thank Alok Patnaik for providing the radio maps used to generate Fig 2 and Lawrence Rudnick for his help. Many thanks to Fred Seward, Paul Plucinsky, and Richard Edgar for suggestions and comments, to Keith Gendreau for his help with detector issues and to the referee, Rob Petre, for his helpful suggestions. IMH would like to thank Ira Glass for his infinite patience and Vivien Marx for her support.

REFERENCES

- Anders, E., & Grevesse, N. 1989, *Geochimica et Cosmochimica Acta*, 53, 197
- Anderson, M. C., & Rudnick, L. 1993, *ApJ*, 408, 514
- Arabadjis, J. S., & Bregman, J. N. 1998, *astro-ph/9806385*
- Blanton, E. L., & Helfand, D. J. 1996, *ApJ*, 470, 961
- Becker, R. H., & Helfand, D. J. 1987, *AJ*, 94, 1629
- Caswell, J. L., Murray, J. D., Roger, R. S., Cole, D. J., & Cooke, D. J. 1975, *A&A*, 45, 239
- Caswell, J. L., Haynes, R. F., Milne, D. K., & Wellington, K. J. 1982, *MNRAS*, 200, 1143
- Condon, J. J. et al 1998, *AJ*, 115, 1693
- Gaensler, B. M. & Johnston, S. 1995, *MNRAS*, 277, 1243
- Ginzburg, V. L. & Syrovatskii, S. I. 1965, *ARA & A*, 3, 297
- Harrus, I. M., Hughes, J. P., & Slane, P. O. 1998, *ApJ*, 499, 273
- Hughes, J. P. & Helfand, D. J., 1985, *ApJ*, 291, 544
- Kaastra, J. S. 1992, *An X-Ray Spectral Code for Optically Thin Plasmas*. Internal SRON-Leiden Report, updated version 2.0
- Mewe, R., Gronenschild, E. H. B. M., & van den Oord, G. H. J. 1985, *A&AS*, 62, 197
- Mewe, R., Lemen, J. R., & van den Oord, G. H. J. 1986, *A&AS*, 65, 511
- Morrison, R. & McCammon, D. 1983, *ApJ*, 270, 119
- Narayan, R. & Schaudt, K. J. 1988, *ApJ*, 325, L43

Patnaik, A. R., Hunt G. C., Salter, C. J., Shaver, P. A. & Velusamy, T. 1990, *A&A*, 232, 467

Raymond, J. C. & Smith, B. W. 1977, *ApJS*, 35, 419

Sedov, L. I. 1959, *Similarity and dimensional methods in mechanics*, (New York: Academic Press)

Seward, F. D. & Wang, Z. R. 1988, *ApJ*, 332, 199

Taylor, G. I. 1950, *Proc Royal Soc London*, 201, 159

Vasisht, G. & Gotthelf, E. V. 1997, *ApJ*, 486L, 129

Fig. 1.— *ASCA* X-ray images of the SNR G39.2–0.3 at low and high energy (top: 0.5–3.0 keV; bottom: $\gtrsim 3.0$ keV) for the GIS (left) and the SIS (right). Contour values are linearly spaced from 30% to 90% of the peak surface brightness in each map. Peak/background values are top: 2.58/0.12; bottom: 1.72/0.08 for the GIS and top: 2.57/0.12; bottom: 1.74/0.19 for the SIS, where all values are quoted in units of $10^{-3}\text{counts s}^{-1} \text{ arcmin}^{-2}$.

Fig. 2.— *Left*: Radio image at 20cm from Patnaik et al (1990) superimposed with SIS high-energy band (from 3.0 to 10.0 keV) contours. The X-ray contours are linearly spaced from 30% to 90% of the peak surface brightness as defined in the Fig. 1. *2.—Right*: Polarized intensity map for data at 6cm with the same X-ray contours superimposed.

Fig. 3.— GIS and SIS spectra – both GIS (GIS 2 and GIS 3) and SIS (SIS 0 and SIS 1) spectra have been combined for display purposes only. The solid curves in the top panels show the joint best-fit NEI thermal plasma plus power-law models. The insert figures show only the thermal component; the bottom panels plot the data/model residuals.

Table 1
Spectral Analysis of Power Law plus NEI Thermal ^a Plasma Model

Parameter	Fit results ^{b,c}
N_H (atoms cm^{-2})	$(4.65^{+0.11}_{-0.26}) \times 10^{22}$
α_p	$2.53^{+0.34}_{-0.27}$
$F_{1\text{keV}}$ (photons $\text{cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1}$)	$(3.4^{+2.5}_{-1.7}) \times 10^{-3}$
Flux ^{α} (ergs $\text{cm}^{-2} \text{ s}^{-1}$) ([0.2 – 3.0] keV)	1.84×10^{-11}
Flux ^{α} (ergs $\text{cm}^{-2} \text{ s}^{-1}$) ([3.0 – 10.0] keV)	2.71×10^{-12}
Flux ^{α} (ergs $\text{cm}^{-2} \text{ s}^{-1}$) ([0.2 – 4.0] keV)	1.92×10^{-11}
kT (keV)	$0.62^{+0.10}_{-0.03}$
$\log(n_e t)$ ($\text{cm}^{-3} \text{ s}$)	$11.08^{+0.19}_{-0.22}$
Normalization ^d (cm^{-5})	$(9.5^{+5.3}_{-3.3}) \times 10^{12}$
Flux ^{Thermal} (ergs $\text{cm}^{-2} \text{ s}^{-1}$) ([0.2 – 3.0] keV)	4.64×10^{-10}
Flux ^{Thermal} (ergs $\text{cm}^{-2} \text{ s}^{-1}$) ([3.0 – 10.0] keV)	7.05×10^{-13}
Flux ^{Thermal} (ergs $\text{cm}^{-2} \text{ s}^{-1}$) ([0.2 – 4.0] keV)	4.65×10^{-10}
Gain _{GIS2} = Gain _{GIS3}	$-3.02^{+0.17}_{-1.54} \%$
χ^2/ν	238.11/235

^a Abundances from Anders & Grevesse (1989)

^b Single-parameter 1 σ errors

^c All fluxes quoted are unabsorbed.

^d $N = (\frac{n_H n_e V}{4\pi D^2})$

Table 2
Summary of the results

Parameter	
R^a (pc)	$11.64 D_{10} \theta_4$
Age (yr)	$6600 - 7300 D_{10} \theta_4$
n_0^b (cm $^{-3}$)	$(0.24^{+0.07}_{-0.05}) D_{10}^{-1/2} \theta_4^{-3/2} f^{-1/2}$
E_0 (ergs)	$(0.35^{+0.15}_{-0.08}) \times 10^{51} D_{10}^{5/2} \theta_4^{3/2} f^{-1/2}$
M_{SU} ($M_\odot$)	$140 - 220 D_{10}^{5/2} f^{1/2}$
P_0 (10^{-10} dyne cm $^{-2}$)	$3.0 - 5.9 D_{10}^{-1/2} f^{-1/2}$
B (μ G)	$90 - 120 D_{10}^{-1/4} f^{-1/4}$
ν_{break} (GHz)	$3900 - 9000 D_{10}^{-5/4} f^{3/4}$
L_X^α (ergs s $^{-1}$) ([0.2 – 4.0] keV)	$2.3 \times 10^{35} D_{10}^2$

^a D_{10} is the distance in units of 10 kpc and θ_4 is the angular radius in units of 4'

^b f is the volume filling factor





